

PRESS RELEASE

Source: Tokyo Institute of Technology

For immediate release: December 22, 2020

Turning the Heat Down: Catalyzing Ammonia Formation at Lower Temperatures with Ruthenium

(Tokyo, December 22) **Scientists at Tokyo Institute of Technology (Tokyo Tech) report that the metal ruthenium, supported with lanthanide oxyhydrides, can efficiently catalyze the synthesis of ammonia at a much lower temperature than the traditional approach. In their new study, they highlight the advantages of the oxyhydride support and its potential in becoming a feasible catalyst for low-temperature ammonia synthesis in the future.**

Nitrogen is an essential nutrient for plant growth. While about 80% of earth is nitrogen, it is mostly contained in the atmosphere as gas, and hence, inaccessible to plants. To boost plant growth, especially in agricultural settings, therefore, chemical nitrogen fertilizers are needed. A crucial step in the production of these fertilizers is the synthesis of ammonia, which involves a reaction between hydrogen and nitrogen in the presence of a catalyst.

Traditionally, ammonia production has been performed through the “Haber-Bosch” process, which, despite being effective, requires high temperature conditions (400–500°C), making the process expensive. Consequently, scientists have been trying to find a way to reduce the reaction temperatures of ammonia synthesis.

Recently, scientists have reported ruthenium—a transition metal—as an efficient “catalyst” for ammonia synthesis, as it operates under milder conditions than traditional iron-based catalysts. However, there is a caveat: nitrogen molecules need to stick to the catalyst surface to undergo dissociation into atoms before reacting with hydrogen to form ammonia. For ruthenium, however, the low temperature often causes hydrogen molecules to stick to its surface instead—a process called hydrogen poisoning—which impedes the production of ammonia. To work with ruthenium, therefore, it is necessary to suppress its hydrogen poisoning.

Fortunately, certain materials can boost the catalytic activity of ruthenium when used as a “catalyst support.” A team of scientists from Tokyo Tech, Japan, recently revealed that lanthanide hydride materials of the form LnH_{2+x} is one such group of support materials. *“The enhanced catalytic performance is realized by two unique properties of the support material. First, they donate electrons, which guide the dissociation of nitrogen on the ruthenium surface. Second, these electrons combine with hydrogen on the surface to form hydride ions, which readily react with nitrogen to form ammonia and release the electrons, suppressing hydrogen poisoning of ruthenium”*, explains Associate Prof. Maasaki Kitano, who led the study.

Suspecting that hydride ion mobility might have a role to play in ammonia synthesis, the team, in a new study published in *Advanced Energy Materials*, investigated the performance of


lanthanide oxyhydrides ($\text{LaH}_{3-2x}\text{O}_x$)—reportedly fast hydride ion conductors at 100-400°C—as a support material for ruthenium, with the aim of uncovering the relationship between ammonia synthesis and hydride ion mobility.

They found that while the “bulk” hydride ion conductivity had little bearing on the activation of ammonia synthesis, the surface or “local” mobility of hydride ions did play a crucial role in catalysis by helping to build up a strong resistance against hydrogen poisoning of ruthenium. They also found that, compared with other support materials, lanthanum oxyhydrides required a lower onset temperature for ammonia formation (160°C) and showed a higher catalytic activity.

Furthermore, the team observed that the presence of oxygen stabilized the oxyhydride framework and the hydride ions against nitridation—the transformation of lanthanum oxyhydride to lanthanum nitride and its subsequent deactivation—which tends to impede catalysis and is a major drawback in using hydride support materials. *“The resistance to nitridation is a tremendous advantage as it helps to preserve the electron donating ability of the hydride ions for longer duration of the reaction,”* comments Prof. Kitano.

The superior catalytic performance and lower synthesis onset temperature achieved using lanthanide oxyhydrides could thus be the much sought-after solution to turn the heat down on ammonia production!

Making Low Temperatures Conducive to Producing Ammonia: Role of Ruthenium Catalysts Promoted by Lanthanide Oxyhydrides



Ruthenium Catalysts Promoted by Lanthanide Oxyhydrides with High Hydride-Ion Mobility for Low-Temperature Ammonia Synthesis
Ooya et al. (2020) | *Advanced Energy Materials*



Reference

Authors: Kayato Ooya¹, Jiang Li¹, Keiga Fukui¹, Soshi Iimura^{1,2}, Takuya Nakao¹, Kiya Ogasawara¹, Masato Sasase¹, Masaaki Kitano^{1,2}, Hitoshi Abe^{3,4,5}, Yasuhiro Niwa³ and Hideo Hosono¹

Title of original paper: Ruthenium Catalysts promoted by Lanthanide Oxyhydrides with High Hydride-Ion Mobility for Low-Temperature Ammonia Synthesis

Journal: *Advanced Energy Materials*
DOI: 10.1002/aenm.202003723
Affiliations: ¹Materials Research Center for Elemental Strategy, Tokyo Institute of Technology
²Precursory Research for Embryonic Science and Technology (PRESTO), Japan
Science and Technology Agency (JST)
³Institute of Materials Structure Science, High Energy Accelerator Research
Organization
⁴Department of Materials Structure Science, School of High Energy Accelerator
Science, SOKENDAI (The Graduate University for Advanced Studies)
⁵Graduate School of Science and Technology, Ibaraki University

*Corresponding author's email: kitano.m.aa@m.titech.ac.jp , hosono@mces.titech.ac.jp

Contact

Kazuhide Hasegawa
Public Relations Group,
Tokyo Institute of Technology
media@jim.titech.ac.jp
+81-3-5734-2975

About Tokyo Institute of Technology

Tokyo Tech stands at the forefront of research and higher education as the leading university for science and technology in Japan. Tokyo Tech researchers excel in fields ranging from materials science to biology, computer science, and physics. Founded in 1881, Tokyo Tech hosts over 10,000 undergraduate and graduate students per year, who develop into scientific leaders and some of the most sought-after engineers in industry. Embodying the Japanese philosophy of “monotsukuri,” meaning “technical ingenuity and innovation,” the Tokyo Tech community strives to contribute to society through high-impact research. <https://www.titech.ac.jp/english/>