

PRESS RELEASE

Source: Tokyo Institute of Technology, Public Relations Group

For immediate release: April 24, 2020

Subject line: **Fuelling the World Sustainably: Synthesizing Ammonia using Less Energy**

(Tokyo, April 24) Scientists at Tokyo Institute of Technology have developed an improved catalyst by taking the common dehydrating agent calcium hydride and adding fluoride to it. The catalyst facilitates the synthesis of ammonia at merely 50 °C, by using only half the energy that existing techniques require. This opens doors to ammonia production with low energy consumption and reduced greenhouse gas emission.




Figure 1. The ammonia challenges

Ammonia (NH_3) is one of the most important industrial chemicals today, synthesized globally for use in fertilizers that then enable food production for approximately 70% of the world's population. Ammonia is currently obtained by reacting nitrogen (N_2) from air with hydrogen (H_2). This reaction requires high energy and is, therefore, powered by fossil fuels, contributing to over 3% of the global CO_2 emissions.

Ammonia is a critical for making plant fertilizer, which in turn feeds approximately 70% of the world's population. In industries, ammonia is produced via the Haber-Bosch process, where methane is first reacted with steam to produce hydrogen, and hydrogen is then reacted with nitrogen to give ammonia. (Figure 1) The problem with this process is that as the temperature increases, the yield decreases. To continue to get a good yield, the pressure applied in the

reaction chamber needs to be increased. This requires much energy. Further, the iron-based catalysts used for the reaction are only effective above 350 °C. Maintaining such high temperatures also requires a significant amount of energy. To top it all, the yield is only 30-40%.

Fossil fuels are currently used to power the process, contributing large amounts of carbon dioxide to the atmosphere. (Figure 1) Renewable resource alternatives, such as wind energy, have been applied, but those have not proven sustainable. To increase the yield while reducing harm to the environment, therefore, the reaction must take place at low temperatures. For this to happen, catalysts that enable the reaction at low temperatures are required.

So far, such catalysts have been elusive to scientists. “Conventional catalysts lose the catalytic activity for ammonia formation from N_2 and H_2 gases at 100-200 °C, even if they exhibit high catalytic performance at high temperatures,” remark a group of scientists from Tokyo Institute of Technology (Tokyo Tech), Japan, who appear to have finally solved the catalyst problem. The scientists, led by Dr. Michikazu Hara, developed a catalyst that is effective even at 50 °C. *“Our catalyst produces ammonia from N_2 and H_2 gases at 50 °C with an extremely small activation energy of 20 kJ mol^{-1} , which is less than half that reported for conventional catalysts,”* Dr. Hara and colleagues report in their paper published in *Nature Communications*.

Their catalyst comprises a solid solution of CaFH, with ruthenium (Ru) nanoparticles deposited on its surface. The addition of fluoride (F^-) to calcium hydride (CaH_2), a common dehydrating agent, is what makes the catalyst effective at lower temperatures and pressures. After conducting spectroscopic and computational analyses, the scientists propose a possible mechanism by which the catalyst facilitates ammonia production.

The calcium–fluoride (Ca–F) bond is stronger than the calcium–hydrogen (Ca–H) bond. So, the presence of the Ca–F bond weakens the Ca–H bond and the Ru is able to extract H atoms from the catalyst crystal, leaving electrons in their place. The H atoms then desorb from the Ru nanoparticles as H_2 gas. This occurs even at 50 °C. The resultant charge repulsion between the trapped electrons and F^- ions in the crystal lower the energy barriers for these electrons to release, thereby giving the material high electron-donating capacity. These released electrons attack the bonds between the nitrogen atoms in the N_2 gas, facilitating the production of ammonia (Figure 2).

This new method of ammonia production cuts energy demands, thereby reducing the carbon dioxide emissions from the use of large amounts of fossil fuels. The findings of this study illuminate the possibility of an environmentally sustainable Haber-Bosch process, opening the door to the next revolution in agricultural food production.




Figure 2. Spectroscopic measurement results and possible reaction mechanism
a) Fourier Transform Infrared (FT-IR) spectra for nitrogen adsorption on the developed fluoride-containing Ru/CaFH catalyst (top) showing weakened bond between nitrogen atoms compared with a non-modified Ru/CaH₂ catalyst (lower spectrum). b) Proposed reaction mechanism using the Ru/CaFH catalyst.


Reference

Authors: Masashi Hattori¹, Shinya Iijima¹, Takuya Nakao², Hideo Hosono², Michikazu Hara^{1,*}
Title of original paper: Solid solution for catalytic ammonia synthesis from nitrogen and hydrogen gases at 50 °C
Journal: *Nature Communications*
DOI: 10.1038/s41467-020-15868-8
Affiliations: ¹Laboratory for Materials and Structures, Tokyo Institute of Technology
²Materials Research Center for Element Strategy, Tokyo Institute of Technology

*Corresponding author's email: mhara@msl.titech.ac.jp, hosono@mces.titech.ac.jp

Videos (Trailer & full versions)

The following videos, provided courtesy of team leader Michikazu Hara, describe the research in a visually and audibly appealing fashion.



A new catalyst to synthesize ammonia from H₂ and N₂ even at 50C (trailer version)

<https://youtu.be/5kmhGJxXaww>



A new catalyst to synthesize ammonia from H₂ and N₂ even at 50C (full version)

<https://youtu.be/91KyC3VgnxQ>

Contact

Emiko Kawaguchi
Public Relations group,
Tokyo Institute of Technology
media@jim.titech.ac.jp
+81-3-5734-2975

About Tokyo Institute of Technology

Tokyo Tech stands at the forefront of research and higher education as the leading university for science and technology in Japan. Tokyo Tech researchers excel in fields ranging from materials science to biology, computer science, and physics. Founded in 1881, Tokyo Tech hosts over 10,000 undergraduate and graduate students per year, who develop into scientific leaders and some of the most sought-after engineers in industry. Embodying the Japanese philosophy of “monotsukuri,” meaning “technical ingenuity and innovation,” the Tokyo Tech community strives to contribute to society through high-impact research.

<https://www.titech.ac.jp/english/>