

PRESS RELEASE

Sources: Tokyo Institute of Technology

For immediate release: April 19, 2019

Subject line: **Photonics: The curious case of the disappearing cylinders**

Study opens a new route to achieving invisibility without using metamaterials

(Tokyo, April 19) **A pair of researchers at Tokyo Institute of Technology describes a way of making a submicron-sized cylinder disappear without using any specialized coating. Their findings could enable invisibility of natural materials at optical frequency^[1] and eventually lead to a simpler way of enhancing optoelectronic devices, including sensing and communication technologies.**

Making objects invisible is no longer the stuff of fantasy but a fast-evolving science. 'Invisibility cloaks' using metamaterials^[2] — engineered materials that can bend rays of light around an object to make it undetectable — now exist, and are beginning to be used to improve the performance of satellite antennas and sensors. Many of the proposed metamaterials, however, only work at limited wavelength ranges such as microwave frequencies.

Now, Kotaro Kajikawa and Yusuke Kobayashi of Tokyo Institute of Technology's Department of Electrical and Electronic Engineering report a way of making a cylinder invisible without a cloak for monochromatic illumination at optical frequency — a broader range of wavelengths including those visible to the human eye.

They firstly explored what happens when a light wave hits an imaginary cylinder with an infinite length. Based on a classical electromagnetic theory called Mie scattering^[3], they visualized the relationship between the light-scattering efficiency of the cylinder and the refractive index^[4]. They looked for a region indicating very low scattering efficiency, which they knew would correspond to the cylinder's invisibility.

After identifying a suitable region, they determined that invisibility would occur when the refractive index of the cylinder ranges from 2.7 to 3.8. Some useful natural materials fall within this range, such as silicon (Si), aluminum arsenide (AlAs) and germanium arsenide (GaAs), which are commonly used in semiconductor technology.

Thus, in contrast to the difficult and costly fabrication procedures often associated with exotic metamaterial coatings, the new approach could provide a much simpler way to achieve invisibility.

The researchers used numerical modeling based on the Finite-Difference Time-Domain (FDTD) method to confirm the conditions for achieving invisibility. (See

Figure 1.) By taking a close look at the magnetic field profiles, they inferred that “the invisibility stems from the cancellation of the dipoles generated in the cylinder.”

Although rigorous calculations of the scattering efficiency have so far only been possible for cylinders and spheres, Kajikawa notes there are plans to test other structures, but these would require much more computing power.

To verify the current findings in practice, it should be relatively easy to perform experiments using tiny cylinders made of silicon and germanium arsenide. Kajikawa says: “We hope to collaborate with research groups who are now focusing on such nanostructures. Then, the next step would be to design novel optical devices.”

Potential optoelectronic applications may include new kinds of detectors and sensors for the medical and aerospace industries.

Image





Figure 1. Computer simulations verify the conditions for achieving invisibility

(a) Light with a wavelength of 700 nm traveling from bottom to top is distorted when the radius of the cylinder (in the middle) is 175 nm. (b) There is hardly any distortion when the cylinder has a radius of 195 nm. These images correspond to the conditions for invisibility predicted by the theoretical calculation.



Yusuke Kobayashi, first author, is a graduate student in the Department of Electrical and Electrical Engineering, School of Engineering at Tokyo Institute of Technology.

Technical terms

[1] Optical frequency: A region of the electromagnetic spectrum that includes ultraviolet, visible and infrared light.

[2] Metamaterials: Specially engineered materials designed to have advantageous electromagnetic properties. Many recent advances in optical cloaking are based on John Pendry's pioneering work on metamaterials.

[3] Mie scattering: A description of the scattering of light particles that have a diameter larger than the wavelength of the incident light. It is named after German physicist Gustav Mie (1868–1957), who published a seminal paper on the scattering of light by gold colloids in 1908.

[4] Refractive index: A measure of how fast light propagates through a material.

References

Yusuke Kobayashi and Kotaro Kajikawa*, Homogeneous dielectric cylinders invisible at optical frequency. *Applied Physics Express* **12**, 042001 (2019), DOI: [10.7567/1882-0786/ab02bb](https://doi.org/10.7567/1882-0786/ab02bb)

Affiliation

Department of Electrical and Electronic Engineering, School of Engineering, Tokyo Institute of Technology, 4259 Nagatsuta, Midori-ku, Yokohama 226-8502, Japan

*Corresponding author's email: kajikawa@ee.e.titech.ac.jp

Related links

Kajikawa research group

<http://www.opt.ip.titech.ac.jp/>

Biometamaterials: Inspiration from the surface of leaves
<https://www.titech.ac.jp/english/news/2015/032670.html>

Controlling Electromagnetic Fields
<https://science.sciencemag.org/content/312/5781/1780>

Invisibility of a finite dielectric cylinder under Fano resonance conditions
<https://link.springer.com/article/10.1134%2FS1063783415100273>

Switchable invisibility of dielectric resonators
<https://journals.aps.org/prb/abstract/10.1103/PhysRevB.95.165119>

Contact

Emiko Kawaguchi
Public Relations Section,
Tokyo Institute of Technology
E-mail: media@jim.titech.ac.jp
+81-3-5734-2975

About Tokyo Institute of Technology

Tokyo Tech stands at the forefront of research and higher education as the leading university for science and technology in Japan. Tokyo Tech researchers excel in fields ranging from materials science to biology, computer science, and physics. Founded in 1881, Tokyo Tech hosts over 10,000 undergraduate and graduate students per year, who develop into scientific leaders and some of the most sought-after engineers in industry. Embodying the Japanese philosophy of “monotsukuri,” meaning “technical ingenuity and innovation,” the Tokyo Tech community strives to contribute to society through high-impact research.

<https://www.titech.ac.jp/english/>