


## PRESS RELEASE

Source: Tokyo Institute of Technology, Public Relations Section  
For immediate release: November 16, 2018

Subject line: **Metallic nano-particles light up another path towards eco-friendly catalysts**  
---Platinum clusters consisting 19 atoms perform 50 times higher catalytic activity---

(Tokyo, November 16) **Scientists at Tokyo Institute of Technology produced subnano-sized metallic particles that are very effective as catalysts for the oxidation of hydrocarbons. These catalysts can be as much as 50 times more effective than well-known Au-Pd bimetallic nanocatalysts.**



**Figure 1. Catalytic activity of subnano-sized metallic particles within dendrimers.**


Each dendrimer molecule hosts a subnano-sized metallic particle that allows for the oxidation of aromatic hydrocarbons, such as toluene (left), to produce useful organic compounds, such as benzoic acid (right). Oxygen molecules are represented in red.

The oxidation of aromatic hydrocarbons is critically important for producing a great variety of useful organic compounds that are used throughout all types of industries. These oxidation processes require the use of catalysts and solvents, which are usually environmentally hazardous. Thus, finding a solvent-free oxidation process using nanosized catalytic particles has attracted considerable attention. Interestingly, sub-nanoscale catalytic particles (subnanocatalysts, or SNCs) composed of noble metals are even better at their job because their increased surface area and unique electronic state results in favorable effects for oxidizing hydrocarbons and also prevents them from getting oxidized themselves. This makes them cost-effective because the amount of metal required for SNCs is lower than for nanosized catalysts.

A team including Dr. Miftakhul Huda, Keigo Minamisawa, Dr. Takamasa Tsukamoto, and Dr. Makoto Tanabe at Tokyo Institute of Technology (Tokyo Tech), led by Prof. Kimihisa Yamamoto, created multiple types of SNCs by using dendrimers, which are tree-like spherical molecules that can be used as a template to contain the desired catalysts. “Dendrimer is


expected to provide internal nanospaces that could be suitable for catalytic conversion in the presence of metal particles,” explains Yamamoto (Figure 1).

The team created various catalysts of different sizes, depending on the noble metal used and the number of atoms of each catalytic particle. They compared their performance to find the best noble metal for making SNCs and then tried to determine the mechanism behind their high catalytic activity. Smaller SNCs were found to be better, while less oxophilic metals (such as platinum) were superior. As shown in Figure 2, the team postulated that the surface of platinum SNCs does not oxidize easily, which makes them reusable and results in the highest catalytic performance of the Pt<sub>19</sub> SNC that can be as high as 50 times more effective than the common Au-Pd nanocatalysts (Figure 3). The team will continue working to shed light on these catalytic phenomena. “The development of a more detailed mechanism including theoretical considerations is currently in progress,” says Tanabe. The applications of such catalysts could greatly contribute for reducing pollution and enhancing our effective use of Earth’s metal resources.



**Figure 2. Mechanism behind the superior catalytic activity of SNCs.**

The larger nanocatalysts and oxophilic SNCs become oxidized on their surface, which makes them less effective as catalysts for the oxidation of hydrocarbons over time. However, less oxophilic SNCs make them very effective and reusable catalysts.



**Figure 3. Subnano particles are more active as catalysts.** (a) Less oxophilic platinum was superior to other noble metals in the aerobic toluene oxidation. (b) The Pt<sub>19</sub> SNC was the highest catalytic performance among other Pt SNCs between 12 and 28 atoms.

#### Related links

[Breakthrough in blending metals: Precise control of multimetallic one-nanometer cluster formation achieved](#)

[How a tetrahedral substance can be more symmetrical than a spherical atom](#)

[A Preparative-Scale Reaction Using Platinum Clusters with a Single-Digit Atomicity Realized](#)

[One-Nanometer Trimetallic Alloy Particles Created](#)

#### Reference

Authors: Miftakhul Huda<sup>1,2</sup>, Keigo Minamisawa<sup>1</sup>, Takamasa Tsukamoto<sup>1,2</sup>, Makoto Tanabe<sup>\*1,2</sup>, and Kimihisa Yamamoto<sup>\*1,2</sup>

Title of original paper: Aerobic Toluene Oxidation Catalyzed by Subnano Metal Particles (Selected as a Very Important Paper (VIP) –less than TOP 5%–)

Journal: *Angewandte Chemie International Edition*

DOI: 10.1002/anie.201809530

Affiliations: <sup>1</sup>Institute of Innovative Research, Tokyo Institute of Technology  
<sup>2</sup>JST–ERATO, Yamamoto Atom Hybrid Project, Tokyo Institute of Technology

\*Corresponding author's email: [yamamoto@res.titech.ac.jp](mailto:yamamoto@res.titech.ac.jp)

#### Contact

Emiko Kawaguchi  
Public Relations Section,  
Tokyo Institute of Technology  
[media@jim.titech.ac.jp](mailto:media@jim.titech.ac.jp)  
+81-3-5734-2975

#### About Tokyo Institute of Technology

Tokyo Tech stands at the forefront of research and higher education as the leading university for science and technology in Japan. Tokyo Tech researchers excel in fields ranging from materials science to biology, computer science, and physics. Founded in 1881, Tokyo Tech hosts over 10,000 undergraduate and graduate students per year, who develop into scientific leaders and some of the most sought-after engineers in industry. Embodying the Japanese philosophy of “monotsukuri,” meaning “technical ingenuity and innovation,” the Tokyo Tech community strives to contribute to society through high-impact research. [www.titech.ac.jp/english/](http://www.titech.ac.jp/english/)