

PRESS RELEASE

Source: Tokyo Institute of Technology, Public Relations Section

For immediate release: October 9, 2018

Headline: **A pheromone-sensing gene that predates land-dwelling vertebrates**

(Tokyo, October 9) **Scientists at Tokyo Institute of Technology have discovered a gene that appears to play a vital role in pheromone sensing. The gene is conserved across fish and mammals and over 400 million years of vertebrate evolution, indicating that the pheromone sensing system is much more ancient than previously believed. This discovery opens new avenues of research into the origin, evolution, and function of pheromone signaling.**

Most land-dwelling vertebrates have both an olfactory organ that detects odors and a vomeronasal organ that detects pheromones, which elicit social and sexual behaviors. It has traditionally been believed that the vomeronasal organ evolved when vertebrates transitioned from living in water to living on land. New research by Masato Nikaido and colleagues at Tokyo Institute of Technology (Tokyo Tech), however, suggests that this organ may be much older than previously believed.

The vomeronasal organ contains receptors in the V1R protein family that are crucial for pheromone detection. Nikaido et al. identified a gene, *ancV1R*, that encodes a previously unknown member of the V1R family. However, unlike other V1R genes, *ancV1R* is present not only in land-dwelling vertebrates but also in some fish lineages, indicating that it has been conserved over 400 million years of vertebrate evolution. According to Nikaido, this finding was “quite surprising, as this represents the first discovery of a V1R family gene shared between fish and mammals.”

The authors identified *ancV1R* in 56 of 115 vertebrate genomes. Interestingly, the loss of *ancV1R* in some vertebrate lineages, such as higher primates (including humans, chimpanzees, and gorillas), cetaceans (including whales and dolphins), birds, and crocodiles, corresponds with the loss of the vomeronasal organ in these lineages (Fig. 1). The findings suggest not only that *ancV1R* may be a vital component of the vomeronasal organ, but also that this organ predates the transition of vertebrates to land, opening a new avenue of research into its origin.

ancV1R is also unusual in that it is expressed in most vomeronasal sensory neurons. In contrast, other V1R proteins follow a “one neuron—one receptor” rule, with only a single receptor being expressed in each neuron. This further demonstrates the importance of *ancV1R* in pheromone sensing. As noted by Nikaido, “It will be fascinating to further investigate how these patterns of expression are regulated and to determine their functional role in chemosensory signaling.”

Reference

Authors: Hikoyu Suzuki^{1,2}, Hidefumi Nishida¹, Hiro Kondo¹, Ryota Yoda¹, Tetsuo Iwata³, Kanako Nakayama¹, Takayuki Enomoto¹, Jiaqi Wu¹, Keiko Moriya-Ito⁴, Masao Miyazaki⁵, Yoshihiro Wakabayashi⁶, Takushi Kishida⁷, Masataka Okabe⁸, Yutaka Suzuki⁹, Takehiko Ito¹, Junji Hirota^{1,3}, Masato Nikaido¹

Title of original paper: A single pheromone receptor gene conserved across 400 million years of vertebrate evolution


Journal: *Molecular Biology and Evolution*

DOI: [10.1093/molbev/msy186](https://doi.org/10.1093/molbev/msy186)

Affiliations: ¹School of Life Science and Technology, Tokyo Institute of Technology
²Nihon BioData Corporation
³Center for Biological Resources and Informatics, Tokyo Institute of Technology
⁴Department of Brain Development and Neural Regeneration, Tokyo Metropolitan Institute of Medical Science
⁵Department of Biological Chemistry and Food Sciences, Faculty of Agriculture, Iwate University
⁶Institute of Livestock and Grassland Science, National Agriculture and Food Research Organization (NARO)
⁷Wildlife Research Center, Kyoto University
⁸Department of Anatomy, The Jikei University of Medicine
⁹Department of Medical Genome Sciences, Graduate School of Frontier Sciences, The University of Tokyo

*Corresponding author's email: mnikaido@bio.titech.ac.jp

A



B




Figure 1. Single emergence and two major losses of *ancV1R* gene during vertebrate evolution.

(A) *ancV1R* emerged in the common ancestor of bony vertebrates (black circle) and was lost in each of the common ancestors (gray circles) of teleost fish and of turtles, alligators, and birds (gray dashed lines and silhouettes). White silhouette and dashed black lines associated with shark indicate that *ancV1R* has not yet emerged.

(B) Examples of *ancV1R* loss events in higher primates (left) and cetaceans (right). The gray branches indicate the lineages in which *ancV1R* has been lost and the vomeronasal organ is absent. The timing of the mutations causing loss of *ancV1R* is shown above or below the gray circles.

Contact

Emiko Kawaguchi
Public Relations Section,
Tokyo Institute of Technology
media@jim.titech.ac.jp
+81-3-5734-2975

About Tokyo Institute of Technology

Tokyo Tech stands at the forefront of research and higher education as the leading university for science and technology in Japan. Tokyo Tech researchers excel in fields ranging from materials science to biology, computer science, and physics. Founded in 1881, Tokyo Tech hosts over 10,000 undergraduate and graduate students per year, who develop into scientific leaders and some of the most sought-after engineers in industry. Embodying the Japanese philosophy of “monotsukuri,” meaning “technical ingenuity and innovation,” the Tokyo Tech community strives to contribute to society through high-impact research. www.titech.ac.jp/english/